

Comparison of Epothilone and Taxol Binding in Yeast Tubulin using Molecular Modeling

Vajihe Akbari ¹, Sharareh Moghim ², and Mohammad Reza Mofid ^{3*}

1. Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Isfahan University of Medical Science, Isfahan, Iran

2. Department of Bacteriology and Virology, Faculty of Medicine, Isfahan University of medical sciences, Isfahan, Iran

3. Department of Biochemistry, School of Pharmacy and Isfahan Pharmaceutical Sciences Research Center, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract

Microtubules are unique cytoskeletal structures that have structural subunits of $\alpha\beta$ tubulin. Taxol is a typical microtubule stabilizing drug. The epothilones are other natural products with similar mechanism of action to taxol. Despite the highly conserved nature of β -tubulin, some organisms like *Saccharomyces cerevesia* (*S.cerevesia*) are resistant to taxol, but sensitive to epothilones. In order to find differences in sensitivity of yeast tubulin to these molecules, we investigated the binding mode of taxol and epothilone A to yeast tubulin using molecular modeling. The multiple sequence alignment of β -tubulin of different species was performed using ClustalW2. Protein structure of yeast β -tubulin was constructed with Swiss Model 8.05 by using 1TVK. Modeled tubulin was superimposed with PyMol on 1JFF for comparison of three-dimensional structure of two proteins. Our results showed that one of the most interesting differences in binding mode of these molecules is residue 227. The His227 in bovine makes a hydrogen bond by means of its δ -nitrogen with epothilone A and by means of its ϵ -nitrogen with taxol. The Asn227 of yeast can play the role of the δ -nitrogen of the imidazole ring of H227, but not of the ϵ -nitrogen of it. So yeast tubulin in contrast to taxol can interact with epothilone A. Due to conservation of essential residues for binding (T274, R282 and Q292), epothilone A in comparison with taxol can tolerate the interchange in the binding pocket (R276I). Our findings may be of a great aid in the rational design of anti-tumor agents that bind to the taxol binding region of tubulin.

Avicenna J Med Biotech 2011; 3(4): 167-175

Keywords: Epothilone A, Microtubules, *Saccharomyces cerevesia*, Taxol, β -tubulin

Introduction

Microtubules are unique cytoskeletal structures that have structural subunits of $\alpha\beta$ tubulin, which are important for intracellular transport and cell division of all eukaryotes ⁽¹⁾. An effective and validated target for tumor chemotherapeutic agents is the microtubule cytoskeleton. These anticancer drugs can be

divided into two main classes: those that inhibit microtubule assembly and those that stabilize microtubules.

Taxol is a typical microtubule stabilizing drug of the first group. Other natural products with the similar mechanism of action to taxol including epothilones, discodermolide, eleu-

therobin, sarcadietyin and laulimalides, have been isolated and are in various stages of pre-clinical/clinical development ⁽²⁾.

In experiment with bovine brain tubulin, epothilones competitively inhibit the binding of taxol to microtubules ^(3,4). This data suggested that there is a common binding site or overlapping site for two compounds, despite having unrelated structures. In comparison with taxol, epothilones are more effective than taxol against some multidrug resistance cell lines and are more water soluble than taxol ⁽³⁾.

Despite the highly conserved nature of β -tubulin across the phyla, organisms show different degrees of susceptibility and resistance to the different groups of antimicrotubule drugs. For example, many protist and fungal tubulins are only slightly affected by taxol ⁽⁵⁾. However, they are structurally normal and can even co-assemble with mammalian tubulins ⁽⁶⁾. It has been reported that sensitivity to this molecule is controlled by relatively few residues. One study ⁽⁷⁾ made a taxol-binding site in yeast β -tubulin, which is normally taxol-resistant, by replacing five residues with their mammalian counterparts (A19K, T23V, G26D, N227H, and Y270F).

According to the high degree of conservation among tubulin proteins and the relative simplicity of the yeast microtubule cytoskeleton, *S.cerevesia* is a suitable organism for studying the microtubule cytoskeleton. Tubulin of the budding yeast *S.cerevesia* shares 76% sequence identity with bovine tubulin, but yeast tubulin isn't sensitive to taxol. In contrast, epothilones can promote the assembly and stabilization of yeast microtubules. So the sequence and structure differ in yeast tubulin and bovine brain tubulin that allow strong binding of epothilones, but not of taxol ⁽⁸⁻¹¹⁾.

In view of the fact that *S.cerevesia* has only one gene (*TUB2*) encoding β -tubulin, this organism was chosen for our study. In our studies, by using molecular modeling we determined the differences in binding interactions with yeast β -tubulin among the micro-

tubule-stabilizing agents, taxol and epothilones. Understanding of the binding mode shared by taxol and other agents with similar taxol-like activity (such as epothilones) will be a great aid in the rational design of anti-tumor agents that bind to the taxol binding region of tubulin.

Materials and Methods

Multiple sequence alignment

Protein sequence alignments and comparisons were performed using a Basic Local Alignment Search Tool (BLAST) program, blastp, against database specification of non-redundant proteins which were available at the National Center for Biotechnology Information (NCBI) Web server, [<http://www.ncbi.nlm.nih.gov/blast/>] ⁽¹²⁾. Amino acid sequences were obtained from NCBI Web server, [<http://www.ncbi.nlm.nih.gov/>]. Multiple sequence alignments were performed using ClustalW2 [<http://www.ebi.ac.uk/clustalw/>], available at the European Bioinformatics Institute (EBI) Web server ⁽¹³⁾. Consensus of amino acid sequence was obtained from Boxshade available at the European Molecular Biology Network (EMBNET) Web server [<http://www.ch.embn.net.org/>].

Known 3D structure of proteins

Three-dimensional protein structures of bovine brain tubulin in complex with taxol (1JFF) and epothiloneA (1TVK) were obtained from the Protein Data Bank (PDB) [<http://www.rcsb.org/pdb/>] ^(14,15).

Molecular modeling

Protein structure of yeast β tubulin was constructed with Swiss Model 8.05 [<http://www.expasy.org/swissmodel/>] ⁽¹⁶⁻¹⁹⁾ by using a bovine brain tubulin in complex with epothiloneA (1TVK). The sequence identity between yeast β tubulin and bovine brain tubulin are 76%. The yeast tubulin structure model was quite similar to the bovine brain tubulin crystal structure. The Root Mean Square Deviation (RMSD) for 1704 polypeptide backbone atoms was 0.11 Å as calculated with Swiss-Pdb Viewer version 4.0.1 [[Downloaded from <http://www.ajmb.org>](http://www.</p>
</div>
<div data-bbox=)

expasy.org/spdbv/]⁽²⁰⁾. We used QMEAN Z-score as an estimate of the model quality. The reliability of the modeled structure was checked by PROCHECK software online.

Molecular modeling visualization and analysis

Modeled yeast tubulin was superimposed with PyMol on the bovine β -tubulin in complex with taxol (1JFF) for comparison of three-dimensional structure of two proteins. Structural presentation of 1JFF and 1TVK and modeled yeast tubulin was made by using Swiss-Pdb Viewer and PyMOL (Version 0.99) [<http://pymol.sourceforge.net>] programs⁽²¹⁾. Cartoon drawings of the structures were obtained using PyMOL⁽²¹⁾. Studies and comparisons of taxol and epothiloneA binding interaction with yeast tubulin were performed by using programs, Swiss-Pdb Viewer and PyMOL.

Results

The multiple sequence alignment of β -tubulin of different species is shown in figure 1. Consensus sequences obtained from the Boxshade program is shown. According to this alignment β -tubulin is highly conserved in different eukaryote.

Tubulin has binding site for GTP that highly conserved among the species. Tubulin like other nucleotide-binding proteins belongs to a class of regulatory proteins called G proteins. Consensus sequence GQC, KGHYTEG, DNEAL and glycine rich sequences SLGGGTGSGMG are binding site for phosphoryl group of nucleotide and are observed in other nucleotide binding proteins (Figure 1)⁽²²⁾. Residues in the 15-27, 212-231, 270-285 and 358-372 region of β -tubulin form hydrophobic sites for taxol⁽²³⁾ and residues in the 212-231 and 270-282 regions make contact with epothilone⁽²⁴⁾.

In these regions, we observed some differences between yeast β -tubulin and β -tubulin of other species. For example, the yeast tubulin instead of K19, V23, D26, H227, A231, F270, R276 and R359 has A, T, G, N, S, Y, I and Q respectively (Figure 1).

The modeled yeast β -tubulin superimposing on the bovine β -tubulin with a RMSD of

0.11 Å and E-value of 4.2e-45 showed that the overall structure is similar (Figure 2). The QMEAN Z-score directly shows how many standard deviations the model's QMEAN score differs from expected values for native or experimental structures (*e.g.* unexpected solvent accessibility, backbone geometry, interatomic packing, *etc.*)⁽²⁵⁾. Good models reach QMEAN Z-scores comparable to experimental structures ($0 < \text{mean Z-score} < 2$).

According to QMEAN Z-score analysis, modeled structure receives a Z-score of -1.8. PROCHECK was used to analyze the Ramachandran plot of the model by comparing it with the Ramachandran plots of 163 representative structures solved at high resolution of at least 2.0 Å. The Ramachandran plot analysis of the model structure showed that 97.20% of the residues are in favor of additional allowed regions. The model was thus evaluated to be of good quality based on these results⁽²⁶⁾. The most striking difference lies in the M loop, which is the taxol binding site. One of the most important residues in M loop that affects binding of taxol to tubulin is residue 276. Comparison of interaction between taxol or epothilone and residue 276 of yeast and mammalian β -tubulin is presented in figure 3.

Analysis of binding interaction of bovine β -tubulin and taxol reveals that 17 residues from the hydrophobic pocket of taxol binding site made direct contact with taxol molecule including V23, D26, L215, L217, H227, L228, A231, S234, F270, P272, L273, T274, S275, R276, P358, R359, G360 (Figure 4A). Seven of these residues are different in yeast β -tubulin (Val23Thr, Asp26Gly, His227Asn, Ala231Ser, Phe270Tyr, Arg276Ile, Arg359Gln replacement) (Figure 4B).

Binding mode analysis of bovine β -tubulin and epothilone A showed that 6 residue from the epothilone A binding site made direct contact with epothilone A molecule including H227, A231, T274, R276, R282, Q292 (Figure 4C). Yeast β -tubulin instead of H227, A231 and R276 has N, S and I respectively (Figure 4D).

Comparison of Epothilone and Taxol




Figure 1. Multiple alignment of β -tubulin of different species: a) *Oryza sativa*, b) *Plasmodium falciparum*, c) *Salmosalar*, d) *Bostaurus*, e) *Drosophila melanogaster*, f) *Homo sapiens*, g) *Cyathostomumcatinatum*, h) *Gallus gallus* and i) *Saccharomyces cerevesia*. Key residues for taxol binding (red), epothilone A binding (orange) and common residues for binding of taxol and epothilone A (green) are highlighted




Figure 2. Superimposition of backbone atoms of modeled yeast β -tubulin (light magenta) on bovine β -tubulin (cyan). Arrows indicates the position of the M loop and residue 276 in the β -tubulin

Molecular modeling shows that another important residue for binding of taxol and epothilone is position 227. Mode of interaction of taxol and epothilone with this residue is compared in figure 5. This comparison shows that these molecules have different mode of interaction with this residue.

Discussion

Our alignment showed the high identity and homology in sequence of β -tubulin of different species. We didn't see any gap except in C terminal. At C-terminal of β -tubulin, we observed extensive molecular heterogeneity. These C terminal regions are highly variable among the isotype within a species




Figure 3. A) Residue 276 of bovine tubulin and its interaction with taxol, B) Residue 276 of yeast tubulin and its interaction with taxol, C) Residue 276 of yeast tubulin and its interaction with epothilone A and D) Residue 276 of bovine tubulin and its interaction with epothilone A

and among species. This region is important for anchoring Microtubule Associated Protein (MAP) in place ⁽²⁷⁾. The diversity of C-terminal may propose the mechanism for isotype-specific MAP binding.

Molecular modeling shows that M loop of yeast β -tubulin in comparison with bovine β -tubulin locates far away taxol molecule. M loop is the main component in contacts between protofilaments, interacting with structures in the H1-S2 and H2-S3 loops of the adjacent monomer ⁽²⁸⁾. *S.cerevesia* exhibit some nonconservative substitutions in the M-loop compared to other species (Figure 1). The interesting substitution is at position 276. The residue R276 in the M loop of bovine β -tubulin, which is a positively charged residue, is replaced by an isoleusine in yeast. This position exists near a predicted bend in the M-loop, and an isoleusine residue here may change the shape of M-loop and its interaction with taxol. Because of the flexible long side chain and its surface location, arginine can interact with taxol.

There are many oxygen groups in taxol molecule that can interact and make hydrogen

bond with positively charged goanidino group of arginine (Figure 3A). This interaction can stabilize and keep the molecule of taxol in its hydrophobic pocket. In contrast, neutral isoleusine cannot make such hydrogen bond and thus cannot stabilize taxol in its pocket binding (Figure 3B). The binding of epothilone A do not affect by this change so the yeast β -tubulin is yet sensitive to this drug. The epothiloneA molecule can interact with yeast tubulin because of existence of important hydrogen bonds between epothilone A and yeast tubulin (Figures 3C and 3D).

The isopropyl group of V23 is near the C3'-benzamidophenyl ring in the bovine β -tubulin (Figure 4A) ⁽²⁹⁾. Yeast β -tubulin has a theronine residue instead of valine. Substitution of non-polar valin by polar threonine leads to distort the positioning of the hydrophobic phenyl ring (Figure 4B). In the bovine β -tubulin, the methylene group of K19, E22, D26 and E27 interact with the C3'-benzamidophenyl ring and stabilize it ⁽²⁸⁾. In yeast β -tubulin, replacement of K19 and D26 by alanin and glycin respectively reduce the methylene groups that interact with the C3'-

Comparison of Epothilone and Taxol




Figure 4. Comparison of the binding mode of paclitaxel and epothilones in yeast and bovine tubulin. Ligands (gray), binding-site key residues (green/cyan) and non-key residues (slate) for bovine tubulin, and, binding-site key residues (salmon) and non-key residues (slate) for yeast tubulin that are shown in stick representation. Hydrogen bonds are presented as yellow long dashes and hydrophobic interactions are shown as red square dots. The remaining part of the β -tubulin secondary structure is rendered with cartoon presentation using PyMOL. A) Taxol as found in the electron crystallography model (PDB: 1JFF). B) Taxol that interacts with modeled yeast tubulin. C) Epothilone A as found in the electron crystallography model (1TVK). D) Epothilone A that interacts with modeled yeast tubulin

benzamido-phenyl ring. The size of epothilone A molecule is smaller than taxol molecule so epothilone A only fill half of taxol binding pocket in β -tubulin. Thus, residues in 15-27 region of β -tubulin including K19, V23 and D26 which interact with taxol, do not play an important role in epothilone binding (Figure 4C)⁽²⁹⁾. We measured distance between hydroxyl group of T23 of yeast β tubulin predicted model and the C3'-benzamido-phenyl ring of taxol molecule, and distance between OH group of T23 of yeast modeled β tubulin and thiazol ring of epothilone. These distances were 5.7 Å for taxol and 9.88 Å for epothilone A. So the polarity of hydroxyl group of threonine can affect the binding of taxol, but not of epothilone A.

In a mutagenesis study, Himes et al reported that epothilones despite taxol can tolerate the change made at position 227 and

bind equally well to wide-type (N227) and the mutated (H227) yeast tubulin⁽⁷⁾.

According to the model, the imidazol ring of H227 of bovine β -tubulin is situated between C2-benzoyl phenyl and C3'-benzamido-phenyl ring of taxol molecule (Figure 4A). The ϵ -nitrogen of imidazol ring of histidine 227 can make hydrogen bond with oxygen atom of C3' benzamido-phenyl ring of taxol, and thus stabilizes this drug in its hydrophobic pocket (Figure 5A). The thiazol ring of epothilone A and the imidazol ring of H227 of bovine β -tubulin are located in a face-to-face orientation (Figure 5B)⁽³⁰⁾. The δ -nitrogen of imidazole ring of H227 can make hydrogen bond with nitrogen atom of thiazol ring of epothilone A (Figure 5B).

In our yeast β -tubulin model, asparagine was situated position 227 (Figure 5A) and because of its distance, δ -nitrogen atom of




Figure 5. Possibility of hydrogen bonding between residue 227 of tubulin [Histidine in bovine (cyan) and Asparagine in yeast (magenta)] and A) oxygen atom of 3'-benzamido group of taxol, B) nitrogen atom of thiazol ring of epothilone A

N227 cannot play the role of ϵ -nitrogen of imidzol ring of H227, thus eliminates the hydrogen bonds (Figure 5A). In contrast, the δ -nitrogen atom of N227 in yeast tubulin can mimic the role of δ -nitrogen atom of histidine and makes hydrogen bond with thiazol ring of epothilone A (Figure 5B).

According to this model, the residue F270 in bovine β -tubulin was located in a hydrophobic basin (Figure 4A) ⁽²⁸⁾. This residue can make van der Waals contact with C3'-phenyl ring and C4- acetyl group of taxol molecule. The epoxide ring at position C12, C13 of epothilone A accommodates the C3'-phenyl ring and C4- acetyl group of taxol ⁽³¹⁾, and is located at the entrance of the hydrophobic basin, but in contrast totaxol, it doesn't enter this pocket. The measured distance between the C3'-phenyl ring of taxol and F270 was 3.8 Å. This distance for epoxide ring of epothilone A and same residue was 8.96 Å. In our yeast β -tubulin model, tyrosine and serine were located at positions 231 and 270. These hydrophilic side chains distort the van der Waals contacts of hydrophobic basin with C3'-phenyl ring and C4- acetyl group of taxol molecule. The residue F270 is a key residue for taxol binding, but not to the epothilone A. So the change of this residue in contrast to taxol does not affect the binding of epothilone to yeast β -tubulin. The cytotoxicity data show that the Phe270Val mutation has only a 3-fold effect on the sensitivity of the epothilone A

compared with a 27-fold change for the taxol ⁽³²⁾, confirm this finding.

The position 359 in bovine β -tubulin is occupied by arginine that makes hydrogen bond with taxol molecule (Figure 4A). In yeast β -tubulin glutamine locates at position 359 and distorts this hydrogen bonding. In agreement with our finding, Henriquez et al reported that sequence replacement of Arg 359Ala in *Acanthamoeba* tubulin lead to resistance to taxol ⁽³³⁾. The distances of this residue with taxol and epothilone A was measured 3.2 and 8.4 Å, respectively. According to this finding, epothilone is so far to make direct hydrogen bound with this residue so that its binding is not affected by changing this residue.

According to the model, the 7-OH group of epothilone and oxetane ring of taxol made hydrogen bonding with T274. The residue T274 is important for binding of both of taxol and epothilone A and it is confirmed by mutagenesis study which reported that a Thr274Ile replacement had a determined effect on activity ⁽³⁴⁾. This effect is more emphasized on epothilone than on taxol activity. This residue is conserved in yeast so the epothilone A can made one of the most important interaction with its β -tubulin.

The model also showed that residue arginine at position 282 is essential for binding of epothilone (Figure 4B), but not of taxol. Again, this residue is conserved in yeast

thereby epothilone A can bind to its β -tubulin. This positively charged residue made electrostatic contacts with the 7-OH group of epothilone A. On the other hand, R282 can form hydrogen binding with T274 and thereby stabilizing conformation of M loop for interaction with epothilone A. In contrast, taxol doesn't make such network of electrostatic interactions and this residue isn't a key residue for its binding and again mutagenesis study agrees with this fact⁽²⁴⁾.

The other residue that is important for binding of epothilone, but not to the taxol is Q292 and again this residue is conserved in yeast, so this organism is sensitive to epothilone A (Figure 4B). This residue locates near the M loop from and forms hydrogen binding with Leu273 adjacent to T274⁽¹⁵⁾. Mutagenesis studies showed that Gln292Glu replacement distorts this hydrogen bonds network of M loop so inhibits epothilone binding⁽³⁴⁾.

Conclusion

Analysis of *S.cerevesia* β -tubulin has shown that failure in interaction with the C3'-benzamidophenyl and phenyl rings and stabilizing them, because of Val23Thr, Asp26Gly, His227Asn, Ala231Ser, Phe270Tyr replacements, is the main reason for resistance to taxol. Study of modeled yeast tubulin revealed that ability of epothilone to stabilize yeast tubulin is related to conservation of key residues for epothilone binding (including T274, R282 and Q292) in this organism. On the other hand, epothilone A can tolerate some interchange in taxol binding pocket residues due to flexibility of epothilone A in comparison with taxol.

Our direct comparison of epothilone binding and taxol binding in yeast tubulin can support development of novel natural and synthetic microtubule stabilizing agents with higher affinity for tubulin and lower sensitivity toward development of resistance of cancerous cells. This would lead to make anticancer drugs with an improved efficiency and fewer side effects.

Acknowledgement

Special thanks are extended to Dr. Ahmad Reza Shahverdi and Dr. Mojtaba Panjehpour for their helpful suggestions. This research was supported by the Bioinformatics Research Center of Isfahan University of Medical Sciences.

References

1. Hyams JS, Lloyd CW. Microtubules. In: Harford JB (eds). Modern Cell Biology. Vol. 13. New York: Wiley-Liss; 1994.
2. He L, Orr GA, Horwitz SB. Novel molecules that interact with microtubules and have functional activity similar to Taxol. Drug Discov Today 2001;6(22):1153-1164.
3. Bollag, DM, McQueney PA, Zhu J, Hensens O, Koupal L, Liesch J, et al. Epothilones, a new class of microtubule-stabilizing agents with a taxol-like mechanism of action. Cancer Res 1995;55(11):2325-2333.
4. Kowalski R J, Giannakakou P, Hamel E. Activities of the microtubule-stabilizing agents epothilones A and B with purified tubulin and in cells resistant to paclitaxel (Taxol[®]). J Biol Chem 1997;272(4):2534-2541.
5. Gull K. Protist tubulins: new arrivals, evolutionary relationships and insights into cytoskeletal function. Curr Opin Microbiol 2001;4(4):427-432.
6. McKean PG, Vaughan S, Gull K. The extended tubulin superfamily. J Cell Sci 2001;114(Pt 15):2723-2733.
7. Gupta ML, Bode CJ, Georg GI, Himes RH. Understanding tubulin-taxol interactions: mutations that impart taxol binding to yeast tubulin. Proc Natl Acad Sci USA 2003;100(11):6394-6397.
8. Little M, Seehaus T. Comparative analysis of tubulin sequences. Comp Biochem Physiol B 1988;90(4):655-670.
9. Burns RG. Alpha-, beta-, and gamma-tubulins: sequence comparisons and structural constraints. Cell Motil Cytoskeleton 1991;20(3):181-189.
10. Brans G, Louie KA, Botstein D. Yeast proteins associated with microtubules in vitro and in vivo. Mol Biol Cell 1992;3(1):29-47.
11. Bode CJ, Gupta ML, Reiff EA, Suprenant KA, Georg GI, Himes RH. Epothilone and paclitaxel: unexpected differences in promoting the assembly and stabilization of yeast microtubules. Biochemistry 2002;41(12):3870-3874.
12. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-

- BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997;25(17):3398-3402.
13. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 1994;22(22):4673-4680.
 14. Lowe J, Li H, Downing KH, Nogales E. Refined structure of alpha beta-Tubulin at 3.5 Å resolution. *J Mol Biol* 2001;313(5):1045-1057.
 15. Nettles JH, Li H, Cornett B, Krahn JM, Snyder JP, Downing KH. The binding mode of epothilone A on alpha, beta-tubulin by electron crystallography. *Science* 2004;305(5685):866-869.
 16. Arnold K, Bordoli L, Kopp J, Schwede T. The SWISS-MODEL Workspace: A web-based environment for protein structure homology modelling. *Bioinformatics* 2006;22(2):195-201.
 17. Kiefer F, Arnold K, Künzli M, Bordoli L, Schwede T. The SWISS-MODEL Repository and associated resources. *Nucleic Acids Res* 2009;37(Database issue):D387-D392.
 18. Schwede T, Kopp J, Guex N, Peitsch MC. SWISS-MODEL: an automated protein homology-modelling server. *Nucleic Acids Res* 2003;31(13):3381-3385.
 19. Guex N, Peitsch MC. SWISS-MODEL and the Swiss-Pdb viewer: an environment for comparative protein modelling. *Electrophoresis* 1997;18(15):2714-2723.
 20. Kabsch W. A discussion of the solution for the best rotation to relate two sets of vectors. *Acta Cryst* 1978;34(Part 5):827-828.
 21. Delano WL. The PyMOL molecular graphics system. DeLano Scientific; San Carlos, CA, USA: 2005.
 22. Sternlicht H, Yaffe MB, Farr GW. A model of the nucleotide-binding site in tubulin. *FEBS Lett* 1987;214(2):226-235.
 23. Xiao H, Verdier-Pinard P, Fernandez-Fuentes N, Burd B, Angeletti R, Fiser A, et al. Insights into the mechanism of microtubule stabilization by Taxol. *Proc Natl Acad Sci USA* 2006;103(27):10166-10173.
 24. Giannakou P, Gussio R, Nogales E, Downing KH, Zaharevitz D, Bollbuck B, et al. A common pharmacophore for epothilone and taxanes: Molecular basis for drug resistance conferred by tubulin mutations in human cancer cells. *Proc Natl Acad Sci USA* 2000;97(6):2904-2909.
 25. Benkert P, Biasini M, Schwede T. Toward the estimation of the absolute quality of individual protein structure models. *Bioinformatics* 2011;27 (3):343-350.
 26. Laskowski RA, MacArthur MW, Moss DS, Thornton JM. PROCHECK: a program to check the stereochemical quality of protein structures. *J Appl Cryst* 1993;26(2):283-291.
 27. Amos LA. Focusing-in on microtubules. *Curr Opin Struct Biol* 2000;10(2):236-241.
 28. Nogales E, Whittaker M, Milligan RA, Downing KH. High-resolution model of the microtubule. *Cell* 1999;96(1):79-88.
 29. Snyder JP, Nettles JH, Cornett B, Downing KH, Nogales E. The binding conformation of Taxol in β -tubulin: A model based on electron crystallographic density. *Proc Natl Acad Sci USA* 2001;98 (9):5312-5316.
 30. Reese M, Sanchez-Pedregal VM, Kubicek K, Meiler J, Blommers MJJ, Griesinger C, et al. Structural basis of the activity of the microtubule-stabilizing agent epothilone a studied by NMR spectroscopy in solution. *Angew Chem Int Ed* 2007;46(11):1864-1868.
 31. Forli S, Manetti F, Altmann KH, Botta M. Evaluation of novel epothilone analogues by means of a common pharmacophore and a QSAR pseudoreceptor model for taxanes and epothilones. *Chem MedChem* 2010;5(1):35-40.
 32. Giannakou P, Sackett DL, Kang YK, Zhan Z, Buters JT, Fojo T, et al. Paclitaxel-resistant human ovarian cancer cells have mutant beta-tubulins that exhibit impaired paclitaxel-driven tubulin polymerization. *J Biol Chem* 1997;272(27):17118-17125.
 33. Henriquez FL, Ingram PR, Muench SP, Rice DW, Roberts CW. Molecular basis for resistance of *Acanthamoeba* tubulins to all major classes of anti-tubulin compounds. *Antimicrob Agents Chemother* 2008;52(3):1133-1135.
 34. He L, Yang CPH, Horwitz SB. Mutations in β -tubulin map to domains involved in regulation of microtubule stability in epothilone-resistant cell lines. *Mol Cancer Ther* 2001;1(1):3-10.