



The Effect of Simulated Physiological Oocyte Maturation (SPOM) and L-Carnitine on Bovine Oocyte Developmental Competence

Ali Malekpour^{1,2}, Abolfazl Shirazi^{1,2*}, Sara Borjian Boroujeni^{1,2}, Ali Sarvari^{1,2}, Mohammad Mehdi Naderi^{1,2}, Mostafa Pournourali^{1,2}, Bahareh Behzadi¹ and Mohammad Mehdi Mehrazar¹

1. Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

2. Sina Fanavaran Mandegar Company, Alborz Science and Technology Park, Kamalshahr, Iran

Abstract

Background: Simulated Physiological Oocyte Maturation (SPOM) mimics *in vitro* the physiological events of oocyte maturation in the presence of cAMP modulators. These modulators increase the intracellular concentrations of cAMP, which inhibits the immediate resumption of meiosis and gives the oocyte more time to gain optimal developmental competence. In addition, L-carnitine helps to increase the energy supply of cells through the β -oxidation of fatty acids. This study aimed to investigate the effect of SPOM and L-carnitine supplementation during *In Vitro Maturation* (IVM) and *In Vitro Culture* (IVC) on the developmental competence of bovine oocytes.

Methods: Ovarian Cumulus Complexes (COCs) were cultured in the presence or absence of forskolin+IBMX during the first 2 hr of IVM (pre-IVM) with or without L-carnitine (LC) during IVM or IVC in six experimental groups as follows: I) pre-IVM (pre-IVM group), II) pre-IVM with L-carnitine supplementation during IVM (pre-IVM/LC group), III) L-carnitine supplementation during IVM (IVM/LC group), IV) L-carnitine supplementation during *in vitro* culture (IVC/LC group), V) pre-IVM+ IVC/LC group, and VI) no treatment during IVM and IVC (Control group). The cleavage and blastocyst rates, the blastocysts' total cells number, and the expression of *Nanog*, *Bax*, *Oct4*, *Cdx2*, and *Ifnt* genes in resulting blastocysts were assessed. To assess differences among experimental groups, a one-way analysis of variance was initially employed, followed by post hoc Fisher LSD. The difference between groups was considered statistically significant when $p < 0.05$.

Results: The cleavage and blastocyst rates in the Pre-IVM and Pre-IVM/LC groups was higher than control group and other groups ($p \leq 0.05$) except for IVC/LC and IVM/LC groups, respectively. The number of blastocyst's Inner Cell Mass (ICM) in pre-IVM and Pre-IVM/LC groups as well as the ratio of ICM/TE were higher than control group ($p < 0.05$). The expression of OCT4, CDX2, and IFNT increased in both the pre-IVM and pre-IVM/LC groups compared to the control group ($p < 0.05$).

Conclusion: In conclusion, the application of SPOM-adapted IVM and L-carnitine during IVM of bovine oocyte improves the quantity and quality of the resulting embryos.

Keywords: 3-Isobutyl-1-methylxanthine, bcl-2-associated X protein, Blastocyst, Carnitine, Cattle, Meiosis, Oocytes, Pre-IVM

To cite this article: Malekpour A, Shirazi A, Borjian Boroujeni S, Sarvari A, Naderi MM, Pournourali M, *et al.* The Effect of Simulated Physiological Oocyte Maturation (SPOM) and L-Carnitine on Bovine Oocyte Developmental Competence. Avicenna J Med Biotech 2024;16(4): 260-267.

Introduction

Livestock play a pivotal role in rural livelihoods and are essential contributors to the economies of developing countries¹. In the last few decades, there has been an increasing focus on using assisted reproduction biotechnologies to improve livestock productivity². Ever since the first calf produced from *In Vitro* Embryo

Production (IVEP) was introduced in 1981, significant progress has been made to enhance the quality of embryo production in bovines³. Despite significant advancements in assisted reproduction biotechnologies, cattle *In Vitro* Production (IVP) still yields blastocyst rates of only 30-40%⁴. This means that the quality of

IVP-produced embryos is generally lower than that of *in vivo* counterparts, especially in terms of cryo-survivability and morphology⁵.

During oocyte maturation, cells acquire the intrinsic competence to support development until the embryo can independently activate its embryonic genome. This is a complex process that is essential for successful embryo development^{6,7}. Successful development of bovine embryos depends on various factors, including the quality of oocytes and sperm, culture media, and the energy source⁸. In the *In Vitro* Maturation (IVM) of the oocyte, there is a time gap between nuclear and cytoplasmic maturation, so that nuclear maturation starts earlier than cytoplasmic maturation. Cytoplasmic maturation of the oocyte, such as the accumulation of mRNA, proteins, redistribution of organelles, and cellular metabolic changes, along with nuclear maturation, are essential for the proper developmental competence of the oocyte. In mammals, cyclic adenosine monophosphate (cAMP) is necessary to maintain meiotic arrest in oocytes. During IVM, when the oocyte is removed from the antral follicle, it experiences a slight reduction in intra-oocyte cAMP concentration, which triggers the resumption of meiosis^{9,10}. There are approaches to delay nuclear maturation by increasing cAMP concentrations before or during IVM to provide an opportunity for cytoplasmic maturation leading to improved oocyte competence and embryonic development¹¹⁻¹³.

In this regard, various physiological and pharmaceutical methods have been used to inhibit the resumption of meiosis in cow oocyte. Physiological methods, such as cultivation in follicular fluid or cultivation in the presence of parietal granulosa cells are among these methods. Regarding pharmacological methods, substances that increase cAMP levels, such as Phosphodiesterase (PDE) inhibitors and Adenylate Cyclase (AC) activators, are effective in inhibiting the resumption of spontaneous meiosis in many animal species^{14,15}.

Forskolin (FSK) and 3-isobutyl-1-methylxanthine (IBMX) are small molecules known to elevate intracellular cAMP concentrations¹⁶. FSK activates the enzyme AC, leading to an increase in intracellular cAMP levels. On the other hand, IBMX acts by blocking non-selective phosphodiesterase activity, preventing the inactivation of intracellular cAMP. This dual action of FSK and IBMX contributes to elevated cAMP concentrations within the cell¹⁷⁻¹⁹. As previously mentioned, the modulation of cAMP levels during IVM plays a crucial role in enhancing the developmental competence of oocytes. Increasing the levels of this second messenger has been shown to significantly improve embryo development²⁰. Therefore, simulating the physiological maturation of the oocyte gives the oocytes more time to undergo appropriate cytoplasmic changes and increases the coordination of the maturation of the nucleus and cytoplasm, and on the other hand, provides

the possibility of having a more homogeneous population of immature oocytes.

L-carnitine is a remarkable compound with several key functions. It serves as a potent antioxidant and is inherently involved in mitochondrial function and lipid metabolism by facilitating the transport of fatty acids to the mitochondria²¹. Moreover, L-carnitine plays a vital role in fatty acid β -oxidation, contributing to increased energy supply to cell²²⁻²⁴. Beyond its metabolic functions, L-carnitine is instrumental in regulating cellular processes, including apoptosis²⁵. It also serves as a guardian of cell membrane integrity and DNA, shielding them from damage caused by Reactive Oxygen Species (ROS).

To evaluate the effect of the presence of cAMP and L-carnitine modulators on the oocyte developmental competence, apart from evaluating the rate of cleavage and blastocyst formation as well as the allocation of blastocyst cells, the expression of some genes related to apoptosis, pluripotency and Maternal Recognition of Pregnancy (MRP) (3 and 43) were evaluated. The studied genes are: *Bax* as pro-apoptotic, *NANOG* and *Oct-4* as markers of ICM and pluripotency, *CDX2* as the primary initiator of trophoblast formation and *IFNT* as an indicator of MRP that inhibits the expression of oxytocin and estrogen receptors. Considering the importance of synchrony between nuclear and cytoplasmic maturation, the aim of this study was to improve the developmental competence of oocytes matured *in vitro*³. To achieve this goal, a protocol was implemented to investigate the possible effects of FSK and IBMX during pre-IVM as well as the effect of L-carnitine supplementation during IVM and *In Vitro* Culture (IVC) in bovine species.

Materials and Methods

Experimental procedures in this study were approved by the Bioethics Committee at Avicenna Research Institute (approval no. 92014). Unless otherwise specified, chemicals were sourced from Sigma Chemical Company.

Oocyte collection and IVM

Upon arrival of slaughterhouse bovine ovaries, the ovaries were washed with normal saline containing antibiotics and any visible follicles with 2-8 mm diameter were aspirated with an 18-gauge needle attached to a 20-ml syringe. Cumulus-Oocyte Complexes (COCs) with more than three layers of cumulus cells and homogenous cytoplasm were selected for experimentation. The COCs were cultured in 50- μ l droplets of maturation media consisting of bicarbonate buffered M199 (475 μ g/ml), NaHCO₃ (105 μ g/ml), sodium pyruvate (2 μ g/ml), penicillin-streptomycin (2.5 μ g/ml), FSH (0.05 IU/ml), LH (0.01 IU/ml), and 10% Fetal Bovine Serum (FBS). During IVM, COCs were cultured for 24 hr at 38.5°C in an atmosphere of 5% CO₂ in air with maximum humidity.

Pre IVM or Simulated Physiological Oocyte Maturation (SPOM)

During Pre IVM, the COCs were cultured for 2 hr in the same culture condition as IVM in hepes buffered TCM-199 supplemented with 0.2 mg/ml Bovine Serum Albumin (BSA), 100 mM sodium pyruvate, 5 mg/ml recombinant human insulin/human transferrin (substantially iron-free), 10,000 IU penicillin, 10 mg/ml streptomycin, 100 μM forskolin, and 500 μM IBMX. The COCs were then cultured in IVM medium for 22 hr at the same culture condition as IVM.

In Vitro Fertilization (IVF)

After maturation the COCs were subjected to IVF medium (10 oocytes/ 50 μl droplet) consisting F-TALP medium, supplemented with 0.2 mM sodium pyruvate, 5 mg/ml BSA (free fatty acid), 20 μg/ml heparin, and 5 μg/ml Penicillin-Streptomycin. The frozen-thawed bovine sperm was subjected to swim-down method using pure sperm gradient (500 μl 40% over 500 μl 80%), 750 g for 10 min. The motile sperm were added to IVF droplets at a final concentration of 2×10⁶ spermatozoa/ml. The oocytes and sperm were co-incubated for 18 hr at 39°C with 5% CO₂ in air and maximum humidity.

In vitro culture of embryos (IVC)

After IVF, the presumptive zygotes were denuded by 3 min vortexing. The zygotes were then cultured in Synthetic Oviductal Fluid (SOF) supplemented with 2% (v/v) BME-essential amino acids, 1% (v/v) MEM-nonessential amino acids, 1 mM glutamine, and 8 mg/ml fatty acid-free BSA. The culture took place at a temperature of 39°C under conditions of 7% O₂, 5% CO₂, and 88% N₂ in humidified air. On the third day of culture, 10% (v/v) charcoal-stripped FBS was added to the medium. The media osmolarity was maintained within the range of 270 to 285 mOsmol/L. The proportions of cleaved embryos, blastocysts, and hatched blastocysts were then determined on days 3, 6-8, and 9, respectively.

Embryo differential staining

To detect Trophectoderm (TE) and Inner Cell Mass (ICM) cells in blastocysts, double staining methods were performed. Briefly, to stain TE cells, blastocysts were exposed to 0.2% Triton X-100 for 20 seconds and then immediately immersed in 30 μg/ml Propidium Iodide (PI) for 1 min. For ICM staining, blastocysts were incubated in ethanol containing 10 μg/ml bisbenzimidazole (Hoechst 33342) for 15 min on ice. Stained blastocysts were mounted on glass slide and examined using an epifluorescence microscope. TE and ICM cells were counted by observing the red and blue colors of the cell nucleus, respectively.

Expression of Bax, Nanog, OCT4, CDX2, and IFNT genes

The expression levels of *Bax*, *Nanog*, *OCT4*, *CDX2*, and *IFNT* genes on day 8, blastocysts of Control, Pre IVM, and Pre IVM/LC groups were determined using quantitative RT-PCR (qRT-PCR). RNA extraction was carried out following the protocol outlined in a previous study³. Subsequently, the concentration of RNA was measured with NanoDrop, and the isolated RNA was stored at -80°C. The Takara cDNA synthesis kit (Takara, Japan, Tokyo) was then utilized to synthesize from the extracted RNA as per the manufacturer's instructions. For relative qRT-PCR analysis, the house-keeping gene was employed as the normalizer gene. The sequences of the primers used for qRT-PCR can be found in table 1. qRT-PCR was conducted using the Step One Plus Real time PCR system (Applied biosystems, USA). Each reaction mixture comprised specific primers, synthesized cDNA, SYBR Green qPCR master mix (Takara, Japan, Tokyo) and distilled water, with reactions set up in triplicates for every run. Three independent biological replicates were performed. The relative qRT-PCR involved an initial denaturation step for 10 min at 95°C, followed by 40 cycles of 15 s at 95°C, 30 s, and 30 s at 72°C. The results were analyzed using the Livak method (2-ΔΔCT formula) within the REST7 program.

Table 1. The primers sequences used in Real Time PCR

| Gene | Sequence of primer (bp) 5' --- 3' | Length (bp) | T _m |
|----------------|---|-------------|----------------|
| Histone H2A FZ | F:TAAAATTGCTGGTGGTGGTGTC R:TTTTACAGAGATACCGTCCACTG | 171 | 60 |
| Oct 4 | F:AGAAAGACGTGGTCCGAGTG R:GCGCCAGAGAAAGGATACG | 138 | 61 |
| CDX2 | F:ACAGTCGCTATATCACCATCC R: TTCCTTTGCTCTGCGGTTCTG | 104 | 64 |
| NANOG | F: TCCTTCTCCTTCAACTCAGCA R: TCCATGGAGGAGGGAAGAGTA | 102 | 60 |
| IFNT | F: AAAGACTCTGACATGGGAAGG R: AAGGTGGTTGATGAAGAGAGG | 155 | 60 |
| BAX | F: TCTGACGGCAACTTCAACTGG R: GAGACAGCCAGGAGAAATCAA | 119 | 64 |

Table 2. The effect of pre-IVM and L-carnitine supplementation on bovine embryo development

| Experimental groups | Oocytes (COCs) n | Blastocyst | | | | | | | | |
|---------------------|------------------|------------------------------|---------|------------|-----------|----------------------------|---------|------------|-----------|------------------------------|
| | | Cleavage n (%) | Day 6 | | | Total n (%) | Day 7 | | | Total n (%) |
| | | | Early n | Expanded n | Hatched n | | Early n | Expanded n | Hatched n | |
| Pre-IVM | 227 | 181 (85.3±3.5) ^a | 28 | 25 | 0 | 53 (30.9±2.4) ^a | 17 | 33 | 25 | 75 (41.2±1.6) ^a |
| Pre-IVM/LC | 130 | 102 (78.3±2.5) ^a | 22 | 7 | 0 | 29 (29.1±3.1) ^a | 14 | 7 | 7 | 38 (43.3±2.1) ^a |
| IVM/LC | 287 | 191 (63.5±3.1) ^b | 34 | 8 | 1 | 43 (22.7±2.6) ^b | 18 | 30 | 23 | 71 (37.9±1.9) ^{a,b} |
| IVC/LC | 89 | 66 (73.0±2.8) ^{a,b} | 18 | 1 | 0 | 19 (28.2±2.4) ^a | 7 | 14 | 1 | 22 (33.4±2.4) ^b |
| Pre IVM+ IVC/LC | 50 | 34 (66.6±3.1) ^b | 7 | 1 | 0 | 8 (23.1±3.2) ^b | 6 | 3 | 2 | 11 (32.3±1.4) ^b |
| Control | 226 | 139 (64.3±3.5) ^b | 17 | 16 | 0 | 33 (23.1±2.0) ^b | 13 | 19 | 12 | 44 (31.5±2.8) ^b |

a,b) Numbers with different lowercase superscript letters in the same column are significantly different ($p < 0.05$).

Experimental groups

In this study, to evaluate the effect of FSK+IBMX and LC on oocyte's developmental competence, the COCs ($n=1009$) were cultured in the presence or absence of Forskolin+IBMX during the first 2 hr of IVM (pre-IVM) with or without L-carnitine (LC) during IVM or IVC. Accordingly, the aspirated COCs were cultured in six experimental groups: I) in the presence of FSK+IBMX during the first 2 hr of IVM (pre-IVM; $n=227$), II) Pre-IVM with L-carnitine supplementation during IVM (pre-IVM/LC group; $n=130$), III) L-carnitine supplementation during IVM (IVM/LC group; $n=287$), IV) L-carnitine supplementation during IVC (IVC/LC group; $n=89$), V) pre-IVM + IVC/LC group ($n=50$), and VI) no supplementation during IVM and IVC (Control group; $n=226$). The study was followed with assessment of blastocysts cells allocation and the expression of *Bax*, *Nanog*, *OCT4*, *CDX2*, and *IFNT* genes in blastocysts of groups I and II, two groups that responded positively to the treatment, compared to the control group.

Statistical analysis

Data collection in this study was conducted across a minimum of four separate replicates. All collected data were presented as the mean value±SEM. To assess differences among the various experimental groups, a one-way analysis of variance (ANOVA) was initially performed, followed by post hoc Fisher LSD using SigmaPlot (version 11.0). In cases where the equal variance test was not met, treatments were compared using the Student-Newman-Keuls Method. Where the normality test failed, the Kruskal-Wallis ANOVA was employed. Differences between groups were deemed statistically significant when the $p \leq 0.05$.

Results

Effects of SPOM-adapted IVM and L-carnitine on bovine embryo development

The cleavage rate in pre-IVM and Pre-IVM/LC groups was higher than the control and other groups




Figure 1. Epifluorescent microscopic imaging of bovine blastocysts produced by different approaches. ICM and TE nuclei were respectively stained with Hoechst 33342 (blue) and PI (red). A) Blastocyst/Pre IVM, B) Blastocyst/Pre IVM group (LC IVM), C) Blastocyst/control (scale bar: 30 μ m).

ICM; Inner Cell Mass, TE; Trophectoderm, and PI; Propidium Iodide.

($p < 0.05$) with exception of the IVC/LC group. The rate of blastocyst on the seventh day in the pre-IVM and pre-IVM/LC groups was higher than the control group and other groups ($p < 0.05$) with exception of the IVM/LC group. There was no significant difference in the cleavage rate and blastocyst rate on the seventh day between the other groups and the control (Table 2).

Effects of SPOM-adapted IVM and L-carnitine on blastocyst cells number

The number of ICM and ratio of ICM/TE in the Pre-IVM and Pre-IVM/LC groups were higher than the control (Figure 1, Table 3; $p < 0.05$).

Effects of SPOM-adapted IVM and L-carnitine on expression of OCT4, CDX2, IFNT, Nanog, and Bax genes

The expression of *OCT4*, *CDX2*, and *IFNT* genes increased in both Pre-IVM and Pre-IVM/LC groups ($p < 0.05$). However, there was no significant difference pre-IVM group and control group in term of *Nanog* and *Bax* gene expression (Figure 2A). Also, the expression of *OCT4* and *IFNT* genes increased in both Pre-IVM and Pre-IVM/LC groups ($p < 0.05$). However, there was no significant difference between pre-IVM and control groups in term of *Nanog*, *CDX2*, and *Bax* gene expression (Figure 2B).

Simulated Physiological Oocyte Maturation

Table 3. Effects of Pre-IVM and Pre-IVM + LC on blastocyst cell number

| Groups | Blastocyst n | Blastocyst cells (mean±SEM) | | | | |
|------------|--------------|-----------------------------|-------------|-------------|-------------------------|-------------------------|
| | | ICM n | TE n | Total n | ICM/TE % | ICM/Total % |
| Pre-IVM | 25 | 30.16±3.39 ^a | 110.76±6.57 | 140.92±7.35 | 27.31±3.39 ^a | 21.40±2.09 ^a |
| Pre-IVM/LC | 22 | 29.12±3.28 ^a | 110.88±5.63 | 140.00±6.53 | 26.32±3.23 ^a | 20.79±2.01 ^a |
| Control | 19 | 16±2.47 ^b | 112.17±5.43 | 128.17±5.55 | 14.31±2.37 ^b | 12.48±1.82 ^b |

a, b) Numbers with different lowercase letters in the same column are significantly different ($p < 0.05$). ICM; Inner Cell Mass, TE; Trophectoderm.




Figure 2. A) The expression of *Bax*, *OCT4*, *CDX2*, *IFNT*, and *Nanog* genes in Pre-IVM and Control groups. B) The expression of *Bax*, *OCT4*, *CDX2*, *IFNT*, and *Nanog* genes in Pre-IVM/LC and control groups. Numbers with different lowercase letters in the same column are significantly different ($p < 0.05$).

Discussion

In addition to culture conditions, embryo quality is directly affected by oocyte developmental competence. The present study specifically investigated two issues in this context: the effect of Simulated Physiological Oocyte Maturation (SPOM) and L-carnitine on the ability of bovine oocytes to develop. These were selected for their potential to enhance oocyte quality during the IVM process. The results of the present study showed that the use of SPOM-adapted IVM system for bovine oocytes can significantly increase the cleavage rate on the third day compared to the control group. In addition, pre-IVM treatment had a significant effect on day 6 and day 7 blastocyst rates. Moreover, in the pre-IVM group, the number of ICM and the ratio of ICM/TE and ICM/total cells were higher than in the control group. As it is known, the improvement of IVM increases the efficiency of the embryo production process, and based on the studies conducted, the increase

in oocyte cAMP levels during the IVM, increases oocyte competence²⁶.

In 2020, Leal *et al*²⁷ conducted a study on cattle to investigate the effectiveness of SPOM-adapted IVM system. They found that the SPOM-adapted IVM system had a positive effect on blastocyst production in cattle without a negative effect on embryo quality. Their findings showed that the SPOM could mimic the physiological events of maturation *in vivo* and could be a useful tool to improve oocyte competence. However, it is important to consider other factors such as the physiological state of the animal, the age of the sample donor, and changes in protocols when interpreting the results of previous studies²⁷. Overall, these findings provide valuable insights into the potential application of the SPOM in the field of reproductive biology²⁷. The results of the present study confirmed the finding of a previous study²⁸ and determined that SPOM-adapted IVM with FSK+IBMX increased the quantity and quality of blastocyst produced²⁸. However, these results differed from those reported by Razza *et al*²⁹, who reported that the use of the current version of the SPOM system may have adverse effects on oocytes and blastocysts and called for optimal protocols to improve oocyte competence.

Moreover, this study showed that the combined use of SPOM-adapted IVM and L-carnitine supplementation in IVM can increase the blastocyst rate on days 6 and 7, as well as the cleavage rate on day 3 compared to control group. Although L-carnitine supplementation in IVC could increase the cleavage and blastocyst rates on the sixth day as much as pre-IVM and pre-IVM/LC groups, but the total blastocyst rate on the seventh day in this group was lower than the two mentioned groups. Instead, when LC was added during IVM, the total blastocyst rate at day 7 was comparable to the pre-IVM and pre-IVM/LC groups (Table 1). On the other hand, the combination of pre-IVM and IVC/LC, did not have any positive effect on both cleavage and blastocyst rates (days 6 and 7) compared to the control group.

Previous studies show controversial results. In some studies, L-carnitine supplementation during IVM, IVF, and IVC increased the oocyte nuclear maturation rate, number of active mitochondria, cleavage, blastocyst, and hatched blastocyst rates^{30,31}, while in other studies,

supplementation of L-carnitine during IVM, IVF, and IVC did not have any positive effect on the above items^{31,32}. For example, Phongnimitr *et al*³¹ reported that while the use of L-carnitine during IVM improved oocyte nuclear maturation, it had no positive effect on development of vitrified oocyte and subsequent embryo development³². Also, the addition of L-carnitine during IVM and IVC before embryo cryopreservation did not have a significant effect on the hatching rate after warming the frozen embryos³¹. Despite these differences of opinion, the general agreement about the positive effect of LC in *in vitro* embryo production in different species is related to its positive effect as a potential antioxidant in reducing the accumulation of ROS and reducing apoptosis. On the other, the reduction of lipid droplets in the oocyte after the use of carnitine also increases the cryotolerance of the embryo^{33,34}.

In this study, although the cleavage and blastocyst rates were higher in the pre-IVM and Pre-IVM/LC groups than in other groups, L-carnitine supplementation in the pre-IVM/LC group could not further increase the above indicators. However, LC supplementation during IVM was able to improve total blastocysts rate to the same extent as the pre-IVM and Pre-IVM/LC groups. Moreover, the addition of LC during the IVC was able to increase the cleavage and day 6 blastocyst rate to the same extent as the pre-IVM and pre-IVM/LC groups (Table 1). In general, considering the accumulation of ROS in oocytes during IVM due to culture *in vitro* and oxidative stress, the use of L-carnitine is beneficial in reducing the amount of ROS during IVM³⁵⁻³⁸.

The use of IBMX and Forskolin at the beginning of oocyte maturation increases the intracellular cAMP level and stops nuclear maturation and therefore maintaining the appropriate cAMP concentration in the oocyte is a vital requirement for the coordination of nuclear and cytoplasmic maturation processes during oocyte maturation^{28,29}. In this study, due to the specific effect of pre-IVM and to some extent L-carnitine on the developmental competence of the oocyte, the quality of blastocysts produced in both the pre-IVM and Pre-IVM/LC groups was higher than the control group, so that the ICM number and ICM/TE ratio were higher in both groups than control group (Table 2; $p < 0.05$).

Regarding gene expression in the present study, *OCT4* expression is essential for maintaining embryonic stem cells pluripotency and ICM functions, as well as for the expression of *Nanog*, *Sox2* and other genes. It can modulate cell fate in the early stages of embryo development and plays an important role in embryo development^{39,40}. *Nanog* is also required to maintain the epiblast pluripotency and commitment in bovine embryo⁴¹. The *CDX2* gene regulates cell lineage specification during bovine early embryonic development, and *IFNT* is responsible for maternal recognition of pregnancy, implantation, and pregnancy maintenance

(3 and 43). In this study, the higher cleavage rate and the quantitative and qualitative increase of produced blastocysts may be related to the significant increase in the expression of *OCT4*, *CDX2* and *IFNT* genes in the pre-IVM and pre-IVM/LC groups compared to the control group.

The reason for the higher expression of *OCT4*, *CDX2* and *IFNT* genes in the pre-IVM and Pre-IVM/LC groups is at least partially due to the better culture conditions in these groups compared to the control group. The improvement of the culture conditions may be basically related to the presence of cAMP modulators; since the addition of L-carnitine to the IVM and IVC media could not improve the positive effect of cAMP modulators on the evaluated indicators, including the expression of *OCT4*, *CDX2*, and *IFNT* genes.

Conclusion

In summary, this study showed that SPOM-adapted IVM with or without L-carnitine significantly increased the developmental competence of bovine oocytes, which was associated with a quantitative and qualitative increase in blastocysts produced. This positive effect, apart from the better synchronization of nuclear and cytoplasmic maturation of the oocyte, may be at least partially related to the increased expression of pluripotency and implantation genes in blastocysts.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Pournourali M, Tarang A, Mashayekhi F. Chromosomal analysis of two buffalo breeds of Mazani and Azeri from Iran. *Iranian Journal of Veterinary Science and Technology* 2015;7(1):22-31.
2. Shirazi A, Naderi MM, Hassanpour H, Heidari M, Borjian S, Sarvari A, Akhondi MM. The effect of ovine oocyte vitrification on expression of subset of genes involved in epigenetic modifications during oocyte maturation and early embryo development. *Theriogenology* 2016;86(9):2136-46.
3. Amini MS, Naderi MM, Shirazi A, Aminafshar M, Boroujeni SB, Pournourali M, et al. Bioactive Materials Derived from Menstrual Blood Stem Cells Enhance the Quality of In Vitro Bovine Embryos. *Avicenna J Med Biotechnol* 2022;14(4):287-93.
4. Naranjo-Gómez JS, Uribe-García HF, Herrera-Sánchez MP, Lozano-Villegas KJ, Rodríguez-Hernández R, Rondón-Barragán IS. Heat stress on cattle embryo: gene regulation and adaptation. *Heliyon* 2021;7(3):e06570.
5. Nejat-Dehkordi S, Ahmadi E, Shirazi A, Nazari H, Shams-Esfandabadi N. Embryo co-culture with bovine amniotic membrane stem cells can enhance the cryosurvival of IVF-derived bovine blastocysts comparable with co-culture with bovine oviduct epithelial cells. *Zygote* 2021;29(2):102-7.

6. Nazari H, Esfandabadi N, Shirazi A, Ahadi AM, Ahmadi E, Safdarinejad A. Effect of Oocyte Maturation Period on Developmental Rate and Sex ratio Distribution of in vitro Produced Bovine Embryos. *Iranian Journal of Applied Animal Science* 2022; 12(2):249-53.
7. Saleh AC, Sabry R, Mastromonaco GF, Favetta LA. BPA and BPS affect the expression of anti-Mullerian hormone (AMH) and its receptor during bovine oocyte maturation and early embryo development. *Reprod Biol Endocrinol* 2021;19(1):119.
8. Turhan A, Pereira MT, Schuler G, Bleul U, Kowalewski MP. Hypoxia-inducible factor (HIF1alpha) inhibition modulates cumulus cell function and affects bovine oocyte maturation in vitro. *Biol Reprod* 2021;104(2):479-91.
9. Abdulkarim A, Balboula A, Badr M, Bedir W, Elmetwalily M, Zaabel S. Comparing in vitro maturation rates in buffalo and cattle oocytes and evaluating the effect of cAMP modulators on maturation and subsequent developmental competence. *Mansoura Veterinary Medical Journal* 2021; 22(3):136-40.
10. Roelen BA. Bovine oocyte maturation: acquisition of developmental competence. *Reprod Fertil Dev* 2020;32(2): 98-103.
11. Hao T, Xu X, Hao H, Du W, Pang Y, Zhao S, et al. Melatonin improves the maturation and developmental ability of bovine oocytes by up-regulating GJA4 to enhance gap junction intercellular communication. *Reprod Fertil Dev* 2021;33(14):760-71.
12. Razza EM, Pedersen HS, Stroebech L, Fontes PK, Kardarmideen HN, Callesen H, et al. Simulated physiological oocyte maturation has side effects on bovine oocytes and embryos. *J Assist Reprod Genet* 2019;36(3):413-24.
13. Yu BY, Subudeng G, Du CG, Liu ZH, Zhao YF, Namei E, Bai Y, Yang BX, Li HJ. Plasminogen activator, tissue type regulates germinal vesicle breakdown and cumulus expansion of bovine cumulus-oocyte complex in vitro. *Biol Reprod* 2019;100(6):1473-81.
14. Carolan, C, Lonergan P, Monget P, Monniaux D, Mermillod P. Effect of follicle size and quality on the ability of follicular fluid to support cytoplasmic maturation of bovine oocytes. *Mol Reprod Dev* 1996;43(4):477-83.
15. Albus FK, Sasseville M, Lane M, Armstrong DT, Thompson JG, Gilchrist RB. Simulated physiological oocyte maturation (SPOM): a novel in vitro maturation system that substantially improves embryo yield and pregnancy outcomes. *Hum Reprod* 2010;25(12): 2999-3011.
16. Raza SH, Abd El-Aziz AH, Abdelnour SA, Easa AA, Alagawany M, Farag MR, et al. The role of forskolin as a lipolytic stimulator during in vitro oocyte maturation and the in vitro embryo production of livestock. *Reprod Domest Anim* 2021;56(12):1486-96.
17. Thongkittidilok C, Doriguzzi N, Nagashima J, Brown M, Chansaenroj A, Songsasen N. Cilostamide and forskolin maintain gap junction function of incubated dog follicles. *Theriogenology* 2020; 142:222-8.
18. Oliveira CS, da Silva Feuchard VL, de Souza Marques SC, Saraiva NZ. Modulation of lipid metabolism through multiple pathways during oocyte maturation and embryo culture in bovine. *Zygote* 2022;30(2):258-66.
19. Zhenwei J, Xianhua Z. Pre-IVM treatment with C-type natriuretic peptide in the presence of cysteamine enhances bovine oocytes antioxidant defense ability and developmental competence in vitro. *Iran J Vet Res* 2019;20(3):173-179.
20. Jasim SA, Ali SA, Fadhil OQ, Rakhmatova MK, Kzar HH, Margiana R, et al. Investigating the Effects of Hydro-alcoholic *Urtica Dioica* Extract and Retinoic Acid on Follicular Development: An Animal Study. *Med J Islam Repub Iran* 2023;37:1.
21. Jiang WJ, Yao XR, Zhao YH, Gao QS, Jin QG, Li YH, XU YN. L-carnitine prevents bovine oocyte aging and promotes subsequent embryonic development. *J Reprod Dev* 2019;65(6):499-506.
22. Jiang W, Li Y, Zhao Y, Gao Q, Jin Q, Yan C, et al. L-carnitine supplementation during in vitro culture regulates oxidative stress in embryos from bovine aged oocytes. *Theriogenology* 2020;143:64-73.
23. Catandi GD, Cheng MH, Chicco AJ, Chen T, Carnevale EM. L-carnitine enhances developmental potential of bovine oocytes matured under high lipid concentrations in vitro. *Anim Reprod Sci* 2023;252:107249.
24. Carrillo-González DF, Hernández-Herrera DY, Maldonado-Estrada JG. The role of L-carnitine in bovine embryo metabolism. A review of the effect of supplementation with a metabolic modulator on in vitro embryo production. *Anim Biotechnol* 2023;34(2):413-23.
25. Chankitisakul V, Somfai T, Inaba Y, Techakumphu M, Nagai T. Supplementation of maturation medium with L-carnitine improves cryo-tolerance of bovine in vitro matured oocytes. *Theriogenology* 2013;79(4):590-8.
26. Demyda-Peyrás S, Dorado J, Hidalgo M, Anter J, De Luca L, Genero E, et al. Effects of oocyte quality, incubation time and maturation environment on the number of chromosomal abnormalities in IVF-derived early bovine embryos. *Reprod Fertil Dev* 2013;25(7):1077-84.
27. Leal GR, Graciosa MAG, Monteiro CAS, Pasolini R, Dos Reis Camargo AJ, Oliveira CS, et al. The SPOM-adapted IVM system improves in vitro production of bovine embryos. *Theriogenology* 2020 Dec;158:277-82.
28. Li HJ, Sutton-McDowall ML, Wang X, Sugimura S, Thompson JG, Gilchrist RB. Extending prematuration with cAMP modulators enhances the cumulus contribution to oocyte antioxidant defense and oocyte quality via gap junctions. *Hum Reprod* 2016;31(4):810-21.
29. Razza EM, Pedersen HS, Stroebech L, Fontes PK, Kardarmideen HN, Callesen H, et al. Simulated physiological oocyte maturation has side effects on bovine oocytes and embryos. *J Assist Reprod Genet* 2019;36(3):413-24.
30. Zhao B, Li H, Zhang H, Ren S, Li Y, Wang X, et al. The effect of L-carnitine supplementation during in vitro maturation on oocyte maturation and somatic cloned embryo development. *Reprod Biol* 2024;24(2):100853.
31. Phongnimitr T, Liang Y, Srirattana K, Panyawai K, Sripunya N, Treetampinich C, et al. Effect of L-carnitine on maturation, cryo-tolerance and embryo developmental

- competence of bovine oocytes. *Anim Sci J* 2013;84(11):719-25.
32. Carrillo-González DF, Rodríguez-Osorio N, Long CR, Vásquez-Araque NA, Maldonado-Estrada JG. L-carnitine supplementation during in vitro maturation and in vitro culture does not affect the survival rates after vitrification and warming but alters Inf-T and ptgs2 gene expression. *Int J Mol Sci* 2020 Aug 5;21(16):5601.
 33. Ghanem N. L-carnitine improved bovine blastocyst rate and quality when supplemented at different preimplantation stages. *Egyptian Journal of Animal Production* 2015; 52(2):89-99.
 34. Shafiei G, Almasi M, Nikzad H, Miyan J, Mahabadi JA, Moshkdanian G. L-carnitine reduces the adverse effects of ROS and up-regulates the expression of implantation related genes in in vitro developed mouse embryos. *Theriogenology* 2020 Mar 15;145:59-66.
 35. Franciosi F, Coticchio G, Lodde V, Tessaro I, Modina SC, Fadini R, et al. Natriuretic peptide precursor C delays meiotic resumption and sustains gap junction-mediated communication in bovine cumulus-enclosed oocytes. *Biol Reprod* 2014 Sep;91(3):61.
 36. Sutton-McDowall ML, Feil D, Robker RL, Thompson JG, Dunning KR. Utilization of endogenous fatty acid stores for energy production in bovine preimplantation embryos. *Theriogenology* 2012;77(8):1632-41.
 37. Wu GQ, Jia BY, Li JJ, Fu XW, Zhou GB, Hou YP, et al. L-carnitine enhances oocyte maturation and development of parthenogenetic embryos in pigs. *Theriogenology* 2011;76(5):785-93.
 38. Simmet K, Zakhartchenko V, Philippou-Massier J, Blum H, Klymiuk N, Wolf E. OCT4/POU5F1 is required for NANOG expression in bovine blastocysts. *Proc Natl Acad Sci USA* 2018 Mar 13;115(11):2770-5.
 39. Kelleher AM, O'Neil E, Benne J, Cecil R, Spencer TE. NANOG is required to form the epiblast and maintain pluripotency in the bovine embryo. *Mol Reprod Dev* 2020;87(1):152-60.
 40. Pournourali M, Tarang A, Farzadi Haghghi S, Yousefi M, Bahadori MH. Polymorphism variant of MnSOD A16V and risk of female infertility in northern Iran. *Taiwan J Obstet Gynecol* 2016;55(6):801-803.
 41. Salimi M, Shirazi A, Norouzian M, Mehrazar MM, Naderi MM, Shokrgozar MA, et al. Histone modifications of H3K4me3, H3K9me3 and lineage gene expressions in chimeric mouse embryo. *Cell J* 2020;22(1):96-105.